

Suitable for:

SMA® ADVANCED Follow-on Milk is tailored for babies from 6 months as part of a varied weaning diet. It contains Zinc and Vitamins A, C & D to help support the normal function of baby's immune system as well as Iron, Vitamin A, Calcium and Omega 3 & 6 at increased levels to meet these infant changing needs for this stage.

SMA® ADVANCED Follow-on Milk contains 2'FL and LNnT which is the latest breakthrough in infant nutrition.

SMA® ADVANCED Follow-on Milk contains 100% whey, partially hydrolysed protein which helps to accelerate gastric emptying¹ and helps with softer stools.²

Not suitable for:

1. Cows' milk protein intolerance/allergy.
2. Lactose intolerance.
3. Inborn errors of metabolism such as phenylketonuria, galactosaemia and galactokinase deficiency.

Shelf life:

SMA® ADVANCED Follow-on Milk powder has a shelf life of 24 months.



800 g

References:
 1. Billeaud C *et al.* Eur J Clin Nutr. 1990; 44: 577-83.
 2. Laura A. Czerkies *et al.* International Journal of Pediatrics, vol. 2018, Article ID 4969576, 7 pages, 2018.

INFORMATION FOR HEALTHCARE PROFESSIONAL USE ONLY

SMA® ADVANCED FOLLOW-ON MILK

From 6 months + data card



Contains 2FL and LNnT

IMPORTANT NOTICE: We believe that breastfeeding is the ideal nutritional start for babies and we fully support the World Health Organization's recommendation of exclusive breastfeeding for the first six months of life followed by the introduction of adequate nutritious complementary foods along with continued breastfeeding up to two years of age. We also recognise that breastfeeding is not always an option for parents. We recommend that healthcare professionals inform parents about the advantages of breastfeeding. If parents choose not to breastfeed, healthcare professionals should inform parents that such a decision can be difficult to reverse and that the introduction of partial bottle-feeding will reduce the supply of breast milk. Parents should consider the social and financial implications of the use of infant formula. As babies grow at different rates, healthcare professionals should advise on the appropriate time for a baby to begin eating complementary foods. Infant formula and complementary foods should always be prepared, used and stored as instructed on the label in order to avoid risks to a baby's health. **SMA® ADVANCED Follow-on Milk** is only suitable for babies over 6 months as part of a mixed diet. It should not be used as a substitute for breast milk during the first 6 months. The decision to start weaning or to use this product before 6 months, should be made only on the advice of a doctor, midwife, health visitor, public health nurse, dietitian or pharmacist, based on baby's individual needs.



UK 0800 081 81 80
www.smahcp.co.uk
 ROI 1800 931 832
www.smahcp.ie

SMA Nutrition, 1 City Place, Gatwick, RH6 OPA
 In the Republic of Ireland: SMA Nutrition, 3030 Lake Drive, Citywest Business Campus, Dublin 24, Ireland



Nutritional information for SMA® ADVANCED Follow-on Milk

	Units	Per 100 ml	Per 100 kcal	Per 100 g powder
Energy		285	419	2139
		kJ	68	100
		kcal		511
Fat	g	3.5	5.1	26
of which, saturates	g	1	1.4	7.3
of which, unsaturates	g	2.2	3.3	16.7
Carbohydrate	g	7.8	11.5	59
of which, sugars	g	7.8	11.5	59
Fibre	g	0.15	0.22	1.14
Protein	g	1.3	1.9	9.7
Salt* (= Sodium x 2.5)	g	0.07	0.1	0.5
Vitamins				
Vitamin A	µg	68	100	510
Vitamin D	µg	0.9	1.4	7
Vitamin E	mg	1.3	2	10
Vitamin K	µg	5.3	7.8	40
Vitamin C	mg	9.2	13.5	69
Thiamin	mg	0.07	0.1	0.5
Riboflavin	mg	0.15	0.23	1.15
Niacin	mg	0.72	1.1	5.4
Vitamin B ₆	mg	0.05	0.07	0.36
Folic acid	µg	10.8	15.9	81
Vitamin B ₁₂	µg	0.15	0.23	1.15
Biotin	µg	1.66	2.45	12.5
Pantothenic acid	mg	0.64	0.94	4.8
Minerals				
Sodium	mg	27	39	200
Potassium	mg	77	114	580
Chloride	mg	51	75	385
Calcium	mg	46	68	347
Phosphorus	mg	27	39	200
Magnesium	mg	6.8	10	51
Iron	mg	0.7	1.04	5.3
Zinc	mg	0.67	1	5
Copper	mg	0.06	0.09	0.4
Manganese	mg	0.01	0.02	0.1
Fluoride	mg	≤0.01	≤0.01	≤0.06
Selenium	µg	2.3	3.3	17
Iodine	µg	10.6	15.7	80
Others				
Taurine	mg	4	6	30
Choline	mg	7.5	11	56
Inositol	mg	4.3	6.3	32
L-Carnitine	mg	1.9	2.8	14
Nucleotides	mg	2	3	15
2'Fucosyllactose	mg	102	149	763
Lacto-N-Neotetraose	mg	51	75	381
Omega 3:				
α-linolenic acid (ALA)	mg	67	98	500
Docosahexaenoic acid (DHA)††	mg	7.6	11.2	57
Omega 6:				
Linoleic acid (LA)	mg	546	802	4100
Arachidonic acid (AA)††	mg	7.6	11.2	57

INFORMATION FOR HEALTHCARE PROFESSIONAL USE ONLY
Theoretical fatty acid profile of SMA® ADVANCED Follow-on Milk

Fatty Acid		Units	Per 100 ml
Saturated			
Caprylic	C8:0	mg	56
Capric	C10:0	mg	62
Lauric	C12:0	mg	378
Myristic	C14:0	mg	141
Palmitic	C16:0	mg	193
Stearic	C18:0	mg	103
Arachidic	C20:0	mg	7
Behenic	C22:0	mg	18
Total saturated		g	0.96
Unsaturated/Monounsaturated			
Palmitoleic	C16:1	mg	2
Oleic	C18:1	mg	1606
Eicosenoic	C20:1	mg	10
Total monounsaturated		g	1.62
Polyunsaturated			
Linoleic	C18:2	mg	546
Linolenic	C18:3	mg	67
Arachidonic	C20:4	mg	7.6
Docosahexaenoic	C22:6	mg	7.6
Total polyunsaturated		g	0.642

Theoretical amino acid profile of SMA® ADVANCED Follow-on Milk

Amino Acid	mg per 100 ml
Essential & Semi-Essential Amino Acids	
Arginine	67
Cystine	38
Histidine	40
Isoleucine	77
Leucine	163
Lysine	135
Methionine	32
Phenylalanine	47
Threonine	80
Tryptophan	28
Tyrosine	61
Valine	71
Other Amino Acids	
Aspartic acid	154
Serine	60
Glutamic acid	235
Proline	67
Glycine	24
Alanine	66

SMA® ADVANCED Follow-on Milk ingredients

Powder (800 g): Lactose (milk), vegetable oils (sunflower, coconut, rapeseed), partially hydrolysed whey protein (milk), 2'fucosyllactose (2'FL), Lacto-N-Neotetraose (LNnT), calcium phosphate, polyunsaturated fatty acids (AA, DHA [fish oil]), magnesium chloride, potassium chloride, L-arginine, potassium phosphate, vitamins (C, pantothenic acid, niacin, E, riboflavin, A, thiamin, B₆, folic acid, K, biotin, D, B₁₂), sodium chloride, L-histidine, choline bitartrate, L-tyrosin, taurine, inositol, ferrous sulphate, L-carnitine, zinc sulphate, antioxidants (tocopherol-rich extract, ascorbyl palmitate), nucleotides (cytidine-, disodium uridine-, adenosine-, disodium guanosine-5'-monophosphate), copper sulphate, manganese sulphate, potassium iodide, sodium selenate.

Scoop size: 4.4 g

Whey:Casein ratio: 100:0

Potential Renal Solute Load: 128 mOsm/l (powder)

Reconstitution rate: 13.3 g powder/100 ml water

Lactose: 59 g/100 g powder

Osmolality: 325 mOsm/kg H₂O (powder)

*Salt is calculated as sodium x 2.5.
Sodium is present for nutritional purposes.
††LCPs = Long Chain Polyunsaturates.